

5.4
#6

$T: \mathbb{P}_2 \rightarrow \mathbb{P}_4$ is defined by $T(\bar{p}(t)) = \bar{p}(t) + t^2 \bar{p}(t)$.

a) If $\bar{p}(t) = 2 - t + t^2$, then

$$T(\bar{p}(t)) = 2 - t + t^2 + 2t^2 - t^3 + t^4$$

$$\Rightarrow \underline{T(\bar{p}(t)) = 2 - t + 3t^2 - t^3 + t^4}$$

b) Take $c \in \mathbb{R}$, $\bar{p}(t) \in \mathbb{P}_2$

$$T(c\bar{p}(t)) = c\bar{p}(t) + ct^2\bar{p}(t)$$

$$= c(\bar{p}(t) + t^2\bar{p}(t))$$

$$= cT(\bar{p}(t))$$

So scalar multiplication is preserved

Take $\bar{p}(t), \bar{q}(t) \in \mathbb{P}_2$

$$T(\bar{p}(t) + \bar{q}(t)) = (\bar{p}(t) + \bar{q}(t)) + t^2(\bar{p}(t) + \bar{q}(t))$$

$$= (\bar{p}(t) + t^2\bar{p}(t)) + (\bar{q}(t) + t^2\bar{q}(t))$$

$$= T(\bar{p}(t)) + T(\bar{q}(t))$$

So vector addition is preserved

$\Rightarrow T$ is a linear transformation

c) $T(1) = 1 + t^2$

$$T(t) = t + t^3$$

$$T(t^2) = t^2 + t^4$$

So the coordinate vectors for the images of $1, t, t^2$ are

$$\begin{bmatrix} 1 \\ 0 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 0 \\ 1 \\ 0 \end{bmatrix}$$

$$\text{So matrix we seek for } T \text{ is } \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{bmatrix}$$

5.4
406

$T: P_3 \rightarrow R^4$ defined by

$$T(\bar{p}) = \begin{bmatrix} \bar{p}(-3) \\ \bar{p}(-1) \\ \bar{p}(1) \\ \bar{p}(3) \end{bmatrix}$$

the matrix we seek is

$$\begin{bmatrix} 1 & -3 & 9 & -27 \\ 1 & -1 & 1 & -1 \\ 1 & 1 & 1 & 1 \\ 1 & 3 & 9 & 27 \end{bmatrix}$$

The columns of the matrix are the images of $1, t, t^2, t^3$ respectively.

7.1
48

The columns of $\begin{bmatrix} \frac{1}{\sqrt{2}} & -\frac{1}{\sqrt{2}} \\ \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} \end{bmatrix}$ form an orthonormal basis for R^2 . So, the matrix is called orthogonal.

Its inverse is $\begin{bmatrix} \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} \\ -\frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} \end{bmatrix}$

7.1
416

$A = \begin{bmatrix} -7 & 24 \\ 24 & 7 \end{bmatrix}$; A's eigenvalues are found as follows:

$$\det \begin{pmatrix} -7-\lambda & 24 \\ 24 & 7-\lambda \end{pmatrix} = (\lambda+7)(\lambda-7) - 24^2 = \lambda^2 - 49 - 576 = 0$$

$(\lambda^2 - 25^2) = 0 \Rightarrow \lambda_1 = -25$ and $\lambda_2 = 25$ are the eigenvalues
The corresponding eigenvectors are $\begin{bmatrix} -4 \\ 3 \end{bmatrix}$ and $\begin{bmatrix} 3 \\ 4 \end{bmatrix}$.

Normalizing them we get an

orthonormal basis the columns of $P =$

$$\text{So, } A = \begin{bmatrix} -\frac{4}{5} & \frac{3}{5} \\ \frac{3}{5} & \frac{4}{5} \end{bmatrix} \begin{bmatrix} -25 & 0 \\ 0 & 25 \end{bmatrix} \begin{bmatrix} -\frac{4}{5} & \frac{3}{5} \\ \frac{3}{5} & \frac{4}{5} \end{bmatrix}$$

$$P = \begin{bmatrix} -\frac{4}{5} & \frac{3}{5} \\ \frac{3}{5} & \frac{4}{5} \end{bmatrix}$$